Functional Ecology 2001
15, 819–822

Author Index

Ackerly, D.D. *see* Anten, N.P.R.
Aizen, M.A. *see* Tadey, M.
Anten, N.P.R. & Ackerly, D.D. Canopy-level photosynthetic compensation after defoliation in a tropical understorey palm, 252
Anten, N.P.R. & Ackerly, D.D. A new method of growth analysis for plants that experience periodic losses of leaf mass, 804
Arnaud, L., Callaghan, A. & Haubruge, E. Insecticide resistance gene transmission by insecticide-susceptible insects, 812
Arnekleiv, J.V. *see* Berg, O.K.
Askew, A.P. *see* Colasanti, R.L.

Baillie, G. *see* Millard, P.
Balaguer, L., Martínez-Ferri, E., Valladares, F., Pérez-Corona, M.E., Baquedano, F.J., Castillo, F.J. & Manrique, E. Population divergence in the plasticity of the response of *Quercus coccifera* to the light environment, 124
Baldwin, I.T. *see* Van Dam, N.M.
Ballabeni, P., Wlodarczyk, M. & Rahier, M. Does enemy-free space for eggs contribute to a leaf beetle's oviposition preference for a nutritionally inferior host plant?, 318
Baquedano, F.J. *see* Balaguer, L.
Bausenwein, U., Millard, P., Thornton, B. & Raven, J.A. Seasonal nitrogen storage in the forb *Rumex acetosa*, 370
Bayley, M. *see* Holmstrup, M.
Bech, C. *see* Berg, O.K.
Berg, O.K., Hendry, A.P., Svendsen, B., Bech, C., Arnekleiv, J.V. & Lohrmann, A. Maternal provisioning of offspring and the use of those resources during ontogeny: variation within and between Atlantic Salmon families, 13
Bessière, J.-M. *see* Grison-Pigé, L.
Björnsson, B.Th. *see* Johnsson, J.I.
Bokma, F. Evolution of body size: limitations of an energetic definition of fitness, 696
Bonal, D. & Guehl, J.-M. Contrasting patterns of leaf water potential and gas exchange responses to drought in seedlings of tropical rainforest species, 490
Boness, D.J. *see* Bowen, W.D.
Bortolotti, G.R. *see* Smits, J.E.
Bouma, T.J., Nielsen, K.L., Van Hal, J. & Koutstaal, B. Root system topology and diameter distribution of species from habitats differing in inundation frequency, 360
Bowen, W.D., Iverson, S.J., Boness, D.J. & Oftedal, O.T. Foraging effort, food intake and lactation performance depend on maternal mass in a small phocid seal, 325
Brodtkorb, E. *see* Jonsson, B.
Brown, G.P. *see* Webb, J.K.
Bryant, D.M. *see* Nudds, R.L.

Cabaret, J. *see* Leignel, V.
Callaghan, A. *see* Arnaud, L.
Campbell, B.D. *see* White, T.A.
Campelo, J. Jr *see* Vourlitis, G.L.
Cannon, K. *see* Wright, I.J.
Carlson, S.L. *see* Meyer, S.E.
Caseiro, F.T. *see* Vourlitis, G.L.
Casper, B.B., Forseth, I.N., Kempenich, H., Seltzer, S. & Xavier, K. Drought prolongs leaf life span in the herbaceous desert perennial *Cryptantha flava*, 740
Castillo, F.J. *see* Balaguer, L.
Chau, A. *see* Mackauer, M.
Clark, P.J., Ewert, M.A. & Nelson, C.E. Physical apertures as constraints on egg size and shape in the Common Musk Turtle, *Sternotherus odoratus*, 70
Colasanti, R.L., Hunt, R. & Askew, A.P. Passage through bird guts causes interspecific differences in seed germination characteristics, 676
Connor, K.F. *see* Hussain, M.
Convey, P. *see* Worland, M.R.
Costa, D.P. *see* Shaffer, S.A.

Dahlgaard, J. *see* Sørensen, J.G.
Daunt, F., Monaghan, P., Wanless, S., Harris, M.P. & Griffiths, R. Sons and daughters: age-specific differences in parental rearing capacities, 211
Davidson, D.L. & Morris, D.W. Density-dependent foraging effort of Deer Mice (*Peromyscus maniculatus*), 575
Dudgeon, S., Kübler, J.E., Wright, W.A., Vadas, R.L. Sr & Petraitis, P.S. Natural variability in zygote dispersal of *Ascophyllum nodosum* at small spatial scales, 595
Dyer, A.R., Goldberg, D.E., Turkington, R. & Sayre, C. Effects of growing conditions and source habitat on plant traits and functional group definition, 85

Ellington, C.P. *see* Hedenström, A.
Ellsworth, D.S. *see* Naumburg, E.
Ericson, L. *see* Strengbom, J.
Espadaler, X. & Gómez, C. Formicine ants comply with the size-grain hypothesis, 136
Ewert, M.A. *see* Clark, P.J.

Farquhar, G.D. *see* Miller, J.M.
Figuerola, J. *see* Negro, J.J.
Fischer, M.K. & Shingleton, A.W. Host plant and ants influence the honeydew sugar composition of aphids, 544
Ford, N.B. *see* Seigel, R.A.
Forseth, I.N. *see* Casper, B.B.
Forseth, T. *see* Jonsson, B.
Fortin, D. An adjustment of the extended contingency model of Farnsworth & Illius (1998), 138

Gagianas, A.A. *see* Tsialtas, J.T.
Garnier, E., Shipley, B., Roumet, C. & Laurent, G. A self-assembling model of resource dynamics and plant growth incorporating plant functional types, 688
Garrido, J. *see* Negro, J.J.
Germino, M.J. & Smith, W.K. Relative importance of microhabitat, plant form and photosynthetic physiology to carbon gain in two alpine herbs, 243
Glöckler, E. *see* Niklaus, P.A.
Goldberg, D.E. *see* Dyer, A.R.
Gómez, C. *see* Espadaler, X.
Grace, J. *see* Meir, P.
Green, A.J. *see* Negro, J.J.
Griffin, K.L., Tissue, D.T., Turnbull, M.H., Schuster, W. & Whitehead, D. Leaf dark respiration as a function of canopy position in *Nothofagus fusca* trees grown at ambient and elevated CO_2 partial pressures for 5 years, 497
Griffiths, R. *see* Daunt, F.
Grison-Pigé, L., Bessière, J.-M., Turlings, T.C.J., Kjellberg, F., Roy, J. & Hossaert-McKey, M. Limited intersex mimicry of floral odour in *Ficus carica*, 551
Grubb, P.J. *see* Sack, L.
Guehl, J.-M. *see* Bonal, D.

Handley, L.L. *see* Tsialtas, J.T.
Harada, Y. *see* Hikosaka, K.
Harris, M.P. *see* Daunt, F.
Haubruge, E. *see* Arnaud, L.
Hayashi, M.M.S. *see* Vourlitis, G.L.
Hedenström, A., Ellington, C.P. & Wolf, T.J. Wing wear, aerodynamics and flight energetics in bumblebees (*Bombus terrestris*): an experimental study, 417
Heibo, E. *see* Magnhagen, C.
Hendry, A.P. *see* Berg, O.K.
Hester, A. *see* Millard, P.

© 2001 British Ecological Society

Higgins, L.E. & Rankin, M.A. Mortality risk of rapid growth in the spider *Nephila clavipes*, 24
Hikosaka, K., Nagashima, H., Harada, Y. & Hirose, T. Implications of geitonogamous pollination for floral longevity in *Iris gracilipes*, 642
Hill, G.E. *see* McGraw, K.J.
Hillman, S.S. *see* Withers, P.C.
Hirose, T. *see* Hikosaka, K.
Hirota, H. *see* Muraoka, H.
Hodge, A. Foraging and the exploitation of soil nutrient patches: in defence of roots, 416
Hoffmann, A.A. *see* Thomson, L.J.
Holanda, J. *see* Vourlitis, G.L.
Holmstrup, M., Sjursen, H., Ravn, H. & Bayley, M. A simple formulation of interaction between individuals competing for light in a monospecific stand, 647
Hossaert-McKey, M. *see* Grison-Pigé, L.
Hunt, R. *see* Colasanti, R.L.
Hussain, M., Kubiske, M.E. & Connor, K.F. Germination of CO_2-enriched *Pinus taeda* L. seeds and subsequent seedling growth responses to CO_2 enrichment, 344

Ingebrigtsen, P.-J. *see* Jonsson, B.
Ishii, H.S. & Sakai, S. Effects of display size and position on individual floral longevity in racemes of *Narthecium asiaticum* (Liliaceae), 396
Ishii, H.S. & Sakai, S. Effects of maternal identity and incubation temperature on Snapping Turtle (*Chelydra serpentina*) growth, 633
Iverson, S.J. *see* Bowen, W.D.

Janzen, F.J. *see* Kolbe, J.J.
Jenni, L. *see* Schaub, M.
Jensen, A.J. *see* Jonsson, B.
Johnson, G.N. *see* Rosevear, M.J.
Johnsson, J.I. & Björnsson, B.Th. Dehydration tolerance and water vapour absorption in two species of soil-dwelling Collembola by accumulation of sugars and polyols, 654
Jonsson, B., Forseth, T., Jensen, A.J. & Næsje, T.F. Thermal performance of juvenile Atlantic Salmon, *Salmo salar* L., 701
Jonsson, B., Jonsson, N., Brodtkorb, E. & Ingebrigtsen, P.-J. Life-history traits of Brown Trout vary with the size of small streams, 310
Jonsson, N. *see* Jonsson, B.

Kassioumi, M.T. *see* Tsialtas, J.T.
Kearney, M. *see* Shine, R.
Kemp, P.D. *see* White, T.A.
Kempenich, H. *see* Casper, B.B.
Kjellberg, F. *see* Grison-Pigé, L.
Kohyama, T. *see* Takenaka, A.
Koizumi, H. *see* Muraoka, H.
Kolbe, J.J. & Janzen, F.J. The influence of propagule size and maternal nest-site selection on survival and behaviour of neonate turtles, 772
Körner, C. *see* Niklaus, P.A.
Korthals, G.W., Smilauer, P., Van Dijk, C. & Van Der Putten, W.H. Linking above- and below-ground biodiversity: abundance and trophic complexity in soil as a response to experimental plant communities on abandoned arable land, 506
Koutstaal, B. *see* Bouma, T.J.
Kropff, M.J. *see* Schippers, P.
Kubiske, M.E. *see* Hussain, M.
Kübler, J.E. *see* Dudgeon, S.
Kvist, A. & Lindström, Å. Basal metabolic rate in migratory waders: intra-individual, intraspecific, interspecific and seasonal variation, 465

Lara, C. & Ornelas, J.F. Nectar 'theft' by hummingbird flower mites and its consequences for seed set in *Moussonia deppeana*, 78
Laurent, G. *see* Garnier, E.
Leathwick, J.R. & Whitehead, D. Soil and atmospheric water deficits and the distribution of New Zealand's indigenous tree species, 233
Leignel, V. & Cabaret, J. Massive use of chemotherapy influences life traits of parasitic nematodes in domestic ruminants, 569
Lind, J. Escape flight in moulting Tree Sparrows (*Passer montanus*), 29
Lindström, Å. *see* Kvist, A.
Loeschcke, V. *see* Sørensen, J.G.
Lohrmann, A. *see* Berg, O.K.
Lopez, O.R. Seed flotation and postflooding germination in tropical *terra firme* and seasonally flooded forest species, 763

McCoull, C.J. *see* Richardson, A.M.M.
McGraw, K.J. & Hill, G.E. Carotenoid access and intraspecific variation in plumage pigmentation in male American Goldfinches (*Carduelis tristis*) and Northern Cardinals (*Cardinalis cardinalis*), 732
Mackauer, M. & Chau, A. Adaptive self superparasitism in a solitary parasitoid wasp: the influence of clutch size on offspring size, 335
Madsen, T. *see* Olsson, M.
Magnhagen, C. & Heibo, E. Gape size allometry in pike reflects variation between lakes in prey availability and relative body depth, 754
Manrique, E. *see* Balaguer, L.
Martínez-Ferri, E. *see* Balaguer, L.
Mas, R.E. *see* Traveset, A.
Matsumoto, J. *see* Muraoka, H.
Meir, P., Grace, J. & Miranda, A.C. Leaf respiration in two tropical rainforests: constraints on physiology by phosphorus, nitrogen and temperature, 378
Metcalfe, N.B. & Steele, G.I. Changing nutritional status causes a shift in the balance of nocturnal to diurnal activity in European Minnows, 304
Meyer, S.E. & Carlson, S.L. Achene mass variation in *Ericameria nauseosus* (Asteraceae) in relation to dispersal ability and seedling fitness, 274
Millard, P., Hester, A., Wendler, R. & Baillie, G. Interspecific defoliation responses of trees depend on sites of winter nitrogen storage, 535; *see also* Bausenwein, U.
Miller, J.M., Williams, R.J. & Farquhar, G.D. Carbon isotope discrimination by a sequence of *Eucalyptus* species along a subcontinental rainfall gradient in Australia, 222
Miranda, A.C. *see* Meir, P.
Moe, S.J., Stenseth, N.C. & Smith, R.H. Effects of a toxicant on population growth rates: sublethal and delayed responses in blowfly populations, 712
Monaghan, P. *see* Daunt, F.
Morris, D.W. *see* Davidson, D.L.
Muraoka, H., Hirota, H., Matsumoto, J., Nishimura, S., Tang, Y., Koizumi, H. & Washitani, I. On the convertibility of different microsite light availabilty indices, relative illuminance and relative photon flux density, 798

Næsje, T.F. *see* Jonsson, B.
Nagashima, H. *see* Hikosaka, K.
Näsholm, T. *see* Strengbom, J.
Naumburg, E., Ellsworth, D.S. & Pearcy, R.W. Crown carbon gain and elevated [CO_2] responses of understorey saplings with differing allometry and architecture, 263
Negro, J.J., Figuerola, J., Garrido, J. & Green, A.J. Fat stores in birds: an overlooked sink for carotenoid pigments?, 297
Nelson, C.E. *see* Clark, P.J.
Nielsen, K.L. *see* Bouma, T.J.
Niklaus, P.A., Glöckler, E., Siegwolf, R. & Körner, C. Carbon allocation in calcareous grassland under elevated CO_2: a combined ^{13}C pulse-labelling/soil physical fractionation study, 43

© 2001 British Ecological Society,

Nishimura, S. *see* Muraoka, H.
Nogueira, J. De S. *see* Vourlitis, G.L.
Nordin, A. *see* Strengbom, J.
Nudds, R.L. & Bryant, D.M. Exercise training lowers the resting metabolic rate of Zebra Finches, *Taeniopygia guttata*, 458

Oftedal, O.T. *see* Bowen, W.D.
Olsson, M. & Madsen, T. Between-year variation in determinants of offspring survival in the Sand Lizard, *Lacerta agilis*, 443
Ornelas, J.F. *see* Lara, C.

Packard, G.C. & Packard, M.J. Environmentally induced variation in size, energy reserves and hydration of hatchling Painted Turtles, *Chrysemys picta*, 481
Packard, M.J. *see* Packard, G.C.
Pärtel, M. & Wilson, S.D. Root and leaf production, mortality and longevity in response to soil heterogeneity, 748
Pearcy, R.W. *see* Naumburg, E.
Pérez-Corona, M.E. *see* Balaguer, L.
Perkins, S.E. & Speakman, J.R. Measuring natural abundance of ^{13}C in respired CO_2: variability and implications for non-invasive dietary analysis, 791
Petersen, M.K. & Sandström, J.P. Outcome of indirect competition between two aphid species mediated by responses in their common host plant, 525
Petraitis, P.S. *see* Dudgeon, S.
Poorter, L. Light-dependent changes in biomass allocation and their importance for growth of rain forest tree species, 113
Priante Filho, N. *see* Vourlitis, G.L.

Rahier, M. *see* Ballabeni, P.
Rankin, M.A. *see* Higgins, L.E.
Raven, J.A. *see* Bausenwein, U.
Ravn, H. *see* Holmstrup, M.
Reich, P.B. *see* Wright, I.J.
Richardson, A.M.M., Swain, R. & McCoull, C.J. Salt spray limits the inland penetration of a coastally restricted invertebrate: a field experiment using landhoppers (Crustacea: Amphipoda: Talitridae), 435
Riera, N. *see* Traveset, A.
Robinson, M. *see* Thomson, L.J.
Rosevear, M.J., Young, A.J. & Johnson, G.N. Growth conditions are more important than species origin in determining leaf pigment content of British plant species, 474
Rothstein, D.E. & Zak, D.R. Photosynthetic adaptation and acclimation to exploit seasonal periods of direct irradiance in three temperate, deciduous-forest herbs, 722
Roumet, C. *see* Garnier, E.
Roy, J. *see* Grison-Pigé, L.
Ryder, J.J. *see* Siva-Jothy, M.T.

Sack, L. & Grubb, P.J. Why do species of woody seedlings change rank in relative growth rate between low and high irradiance?, 145
Sadras, V.O. *see* Thomas, H.
Sakai, S. *see* Ishii, H.S.
Sandström, J.P. *see* Petersen, M.K.
Sayre, C. *see* Dyer, A.R.
Schaub, M. & Jenni, L. Variation of fuelling rates among sites, days and individuals in migrating passerine birds, 584
Schippers, P. & Kropff, M.J. Competition for light and nitrogen among grassland species: a simulation analysis, 155
Schuster, W. *see* Griffin, K.L.
Seal, N.J. *see* Stockley, P.
Seigel, R.A. & Ford, N.B. Phenotypic plasticity in reproductive traits: geographical variation in plasticity in a viviparous snake, 36
Seltzer, S. *see* Casper, B.B.
Shaffer, S.A., Weimerskirch, H. & Costa, D.P. Functional significance of sexual dimorphism in Wandering Albatrosses, *Diomedea exulans*, 203
Shine, R. & Kearney, M. Field studies of reptile thermoregulation: how well do physical models predict operative temperatures?, 282; *see also* Webb, J.K.
Shingleton, A.W. *see* Fischer, M.K.
Shipley, B. *see* Garnier, E.
Siegwolf, R. *see* Niklaus, P.A.
Siva-Jothy, M.T. & Ryder, J.J. Assaying PHA-induced mitosis: out of control?, 813
Sjursen, H. *see* Holmstrup, M.
Smilauer, P. *see* Korthals, G.W.
Smit, A.-M. & Van Aarde, R.J. The influence of millipedes on selected soil elements: a microcosm study on three species occurring on coastal sand dunes, 51
Smith, R.H. *see* Moe, S.J.
Smith, W.K. *see* Germino, M.J.
Smits, J.E., Bortolotti, G.R. & Tella, J.L. Measurement repeatability and the use of controls in PHA assays: a reply to Siva-Jothy & Ryder, 815
Sørensen, J.G., Dahlgaard, J. & Loeschcke, V. Genetic variation in thermal tolerance among natural populations of *Drosophila buzzatii*: down regulation of Hsp70 expression and variation in heat stress resistance traits, 289
Speakman, J.R. *see* Perkins, S.E.
Spotila, J.R. *see* Steyermark, A.C.
Steele, G.I. *see* Metcalfe, N.B.
Stenseth, N.C. *see* Moe, S.J.
Sterner, R.W. *see* Urabe, J.
Steyermark, A.C. & Spotila, J.R. Asymmetric competition between plant species, 624
Stockley, P. & Seal, N.J. Plasticity in reproductive effort of male dung flies (*Scatophaga stercoraria*) as a response to larval density, 96
Strengbom, J., Nordin, A., Näsholm, T. & Ericson, L. Slow recovery of boreal forest ecosystem following decreased nitrogen input, 451
Svendsen, B. *see* Berg, O.K.
Swain, R. *see* Richardson, A.M.M.

Tadey, M. & Aizen, M.A. Why do flowers of a hummingbird-pollinated mistletoe face down?, 782
Takahashi, K. *see* Takenaka, A.
Takenaka, A., Takahashi, K. & Kohyama, T. Growth-enhanced fish can be competitive in the wild, 660
Tang, Y. *see* Muraoka, H.
Tella, J.L. *see* Smits, J.E.
Thomas, H. & Sadras, V.O. The capture and gratuitous disposal of resources by plants, 3
Thomson, L.J., Robinson, M. & Hoffmann, A.A. Field and laboratory evidence for acclimation without costs in an egg parasitoid, 217
Thornton, B. *see* Bausenwein, U.
Tissue, D.T. *see* Griffin, K.L.
Traveset, A., Riera, N. & Mas, R.E. Optimal leaf display and biomass partitioning for efficient light capture in an understorey palm, *Licuala arbuscula*, 669
Tsialtas, J.T., Handley, L.L., Kassioumi, M.T., Veresoglou, D.S. & Gagianas, A.A. Interspecific variation in potential water-use efficiency and its relation to plant species abundance in a water-limited grassland, 605
Turkington, R. *see* Dyer, A.R.
Turlings, T.C.J. *see* Grison-Pigé, L.
Turnbull, M.H. *see* Griffin, K.L.

Urabe, J. & Sterner, R.W. Contrasting effects of different types of resource depletion on life-history traits in *Daphnia*, 165

Vadas, R.L. *see* Dudgeon, S.
Valladares, F. *see* Balaguer, L.

© 2001 British Ecological Society,

Van Aarde, R.J. *see* Smit, A.-M.
Van Dam, N.M. & Baldwin, I.T. Competition mediates costs of jasmonate-induced defences, nitrogen acquisition and transgenerational plasticity in *Nicotiana attenuata*, 406
Van Damme, R. & Vanhooydonck, B. Origins of interspecific variation in lizard sprint capacity, 186
Van Der Putten, W.H. *see* Korthals, G.W.
Van Dijk, C. *see* Korthals, G.W.
Van Hal, J. *see* Bouma, T.J.
Vanhooydonck, B. *see* Van Damme, R.
Veresoglou, D.S. *see* Tsialtas, J.T.
Vourlitis, G.L., Priante Filho, N., Hayashi, M.M.S., Nogueira, J. De S., Caseiro, F.T., Holanda, J. & Campelo, J. Jr. Seasonal variations in the net ecosystem CO_2 exchange of a mature Amazonian transitional tropical forest (cerradão), 388

Wanless, S. *see* Daunt, F.
Washitani, I. *see* Muraoka, H.
Webb, J.K., Brown, G.P. & Shine, R. Body size, locomotor speed and antipredator behaviour in a tropical snake (*Tropidonophis mairii*, Colubridae): the influence of incubation environments and genetic factors, 561
Weimerskirch, H. *see* Shaffer, S.A.
Wendler, R. *see* Millard, P.
Westoby, M. *see* Wright, I.J.
White, T.A., Campbell, B.D. & Kemp, P.D. Laboratory screening of juvenile responses of grassland species to warm temperature pulses and water deficits to predict invasiveness, 103
Whitehead, D. *see* Griffin, K.L. *and* Leathwick, J.R.
Williams, J.B. Energy expenditure and water flux of free-living Dune Larks in the Namib: a test of the reallocation hypothesis on a desert bird, 175
Williams, R.J. *see* Miller, J.M.
Wilson, S.D. *see* Pärtel, M.
Withers, P.C. & Hillman, S.S. Allometric and ecological relationships of ventricle and liver mass in anuran amphibians, 60
Wlodarczyk, M. *see* Ballabeni, P.
Wolf, T.J. *see* Hedenström, A.
Worland, M.R. & Convey, P. Rapid cold hardening in Antarctic microarthropods, 515
Wright, I.J. & Cannon, K. Relationships between leaf lifespan and structural defences in a low-nutrient, sclerophyll flora, 351
Wright, I.J., Reich, P.B. & Westoby, M. Strategy shifts in leaf physiology, structure and nutrient content between species of high- and low-rainfall and high- and low-nutrient habitats, 423
Wright, W.A. *see* Dudgeon, S.

Xavier, K. *see* Casper, B.B.

Young, A.J. *see* Rosevear, M.J.

Zak, D.R. *see* Rothstein, D.E.

© 2001 British
Ecological Society,

Functional Ecology 2001
15, 823–826

Subject Index

abscission 3–12
acclimation 217–221, 722–731
Acer rubrum 263–273
Achillea millefolium 605–614
adaptation 474–480, 701–711, 722–731
Adenostyles alliariae 318–324
aerodynamics 417–422
air saturation deficit 233–242
Alaskozetes antarcticus 515–524
algae 165–174, 595–604
Allium tricoccum 722–731
allocation 113–123, 406–415, 660–668
allometry 60–69, 263–273, 465–473, 754–762
alpine snowbank 243–251
amino acids 60–69, 370–377, 525–534
analyses 85–95, 113–123, 155–164, 804–811
annual species 85–95, 360–369, 642–646
Anser anser 297–303
ant attendance 544–550
Antarctic 515–524
anthelmintic 569–574
Anthoxanthum odoratum 155–164
aphids 335–343, 525–534, 544–550
Ascophyllum nodosum 595–604
Asparagus 669–675
asymmetric competition 155–164, 615–623
Austrotroides maritimus 435–442

basal metabolic rate 175–185, 465–473
beetles 318–324
Betula pendula 535–543
biodiversity 506–514, 676–687
biomass allocation 804–811
biomass partioning 113–123, 660–668
biophysics 282–288
birds 78–84, 669–675, 782–790:
  desert 175–185
  migratory 297–303, 465–473, 584–594
  passeriformes 29–35, 458–464, 584–594, 732–739
  sea 203–210, 211–216
blooming schedule 633–641
blowfly 712–721
Bombus terrestris 413–422
Brachythecium reflexum 451–457
breath samples 791–797
Brown Trout 310–317, 654–659
bryophytes 451–457
bumblebees 417–422
burrowing 60–69

cadmium 712–721
Caltha leptosepala 243–251
canopy 252–262, 497–505, 642–646
carbohydrates 497–505
carbon 3–12, 49–50, 165–174:
  cycling 388–395
  gain 243–251, 252–262, 263–273, 423–434
  isotope discrimination 222–232, 605–614, 791–797
  nitrogen : phosphorus ratio 165–174
carbon dioxide 388–395, 791–797:
  elevated 43–50, 263–273, 344–350, 497–505
Cardinalis cardinalis 732–739
Carduelis tristis 732–739
carotenoids 297–303, 474–480, 732–739
Carya illinoensis 525–534
Centrobolus spp. 51–59
Chaitophorus spp. 544–550
Chamaedorea elegans 252–262, 804–811
Chelydra serpentina 624–632, 772–781
chemotherapy size increase 569–574
Chenopodiaceae 360–369
chlorophyll 474–480, 722–731
Chrysemys picta 481–489
Cirsium spinosissimum 318–324
coastal sand dunes 51–59
coastal zone 435–442
cold tolerance 515–524
collembola 515–524, 647–653
coloration of birds 297–303, 732–739
competition 575–583, 605–614, 654–659, 676–687:
  larval 712–721
  light 155–164, 252–262, 642–646
competitive asymmetry 155–164, 615–623
contest competition 615–623
copper models 282–288
costs 24–28, 78–84, 396–405, 406–415,
  flight 417–422
critical load 451–457
cross-over point irradiance 145–154
crown architecture 263–273, 660–668
Crozet Archipelago 203–210
Cryptantha flava 740–747
Cryptopygus antarcticus 515–524

daily variations 688–695
Daphnia obtusa 165–174
defoliation 804–811
dehydration tolerance 60–69
desert 175–185, 740–747:
  and Mediterranean 186–202
desiccation tolerance 103–112, 647–653
dicots 85–95, 360–369
diel rhythm 304–309
dietary analysis 791–797
diffuse site factor 798–803
Diomedea exulans 203–210
Diplotropis purpurea 490–496
dispersal 274–281, 595–604, 647–653, 763–771
display size 396–405, 633–641
distributional limits 435–442
Drosophila buzzatti 289–296
drought 740–747:
  soil and atmosphere 233–242, 490–496
dung flies 96–102

ecomorphology 186–202
ecophysiology 388–395
ecosystem recovery 451–457
ecotoxicity 712–721
ecotypes 124–135
ectotherm 282–288
egg 13–23, 70–77, 318–324:
  parasitoid 217–221, 335–343
  variation 481–489
ejaculate expenditure 96–102, 782–790
embryo 481–489
energetics 175–185, 325–334, 417–422
energy expenditure 458–464
energy reserves 304–309
environmental change 85–95
environmental gradients 676–687
environmental variation 481–489
Eperua falcata 490–496
Ericameria nauseosus 274–281
Erythronium grandiflorum 243–251
escape flight 29–35, 417–422
Eucalyptus spp. 222–232
evergreen and deciduous trees 535–543
evolution 3–12, 186–202
evolutionary stable strategy 642–646

© 2001 British Ecological Society

fall rate 274–281
fat stores 13–23, 297–303
fattening rate 584–594
fecundity 36–42, 569–574, 712–721
female reproductive success 443–450, 782–790
Festuca ovina 155–164
Festuca rubra 748–753
Festuca valida 605–614
Ficus carica 551–558
field metabolic rate 175–185
figs 551–558
fish 304–309, 754–762:
  Salmo spp. 13–23, 310–317, 654–659, 701–711
fitness 24–28, 78–84, 335–343, 406–415
flight 29–35, 203–210, 417–422, 458–464
flooding 360–369, 763–771
floral longevity 396–405, 633–641
floral odour mimicry 551–558
flower orientation 782–790
fluctuating selection 443–450
Folsomia fimetaria 647–653
food consumption 701–711
foraging 24–28, 203–210, 325–334, 575–583
forbs 370–377, 605–614
forest 145–154, 263–273, 344–350:
  boreal 451–457, 575–583
  cloud 78–84
  Mediterranean 124–135
  temperate 232–242, 535–543, 722–731, 782–790
  tropical rain 113–123, 252–262, 378–387, 490–496
    and upland 763–771
  tropical transitional 388–395
  understorey 263–273, 798–803
    palms 252–262, 660–668, 804–811
frogs 60–69
frugivory 669–675
fruitfly 289–296
functional classifications 85–95, 676–687, 688–695

gape size variation 754–762
gas exchange responses 43–56, 490–496
generalized additive models 233–242
genetic variability 124–135, 289–296
geographic variation 36–42
germination 103–112, 344–350, 406–415, 669–675, 763–771
giving-up density 575–583
grassland 85–95, 103–112, 370–377:
  calcareous 43–50
  perennials 155–164, 748–753
  semi-arid 605–614
growth 211–216:
  analysis 113–123, 804–811
  conditions 474–480
  efficiency 701–711
  hormone 654–659
  rates 24–28, 624–632, 654–659, 712–721
gut clearance 515–524

halophytic species 360–369
Halozetes belgicae 515–524
headspace 551–558
heat shock proteins 289–296
herbaceous 722–731, 740–747
herbivory 165–174, 406–415, 535–543, 804–811
herbs 243–251, 669–675, 722–731
Holcus lanatus 155–164
honeydew quality 544–550
host range evolution 318–324
hummingbirds 78–84, 782–790
hydrochory 763–771
Hylocomium splendens 451–457

incubation 175–185, 481–489, 772–781:
  temperature 561–568, 624–632
individual-based automata 676–687
inflorescence 78–84, 396–405, 633–641, 782–790
internal cycling 370–377, 535–543
intersex mimicry 551–558
interspecific competition 155–164, 525–534
interspecific defoliation 535–543
interspecific variation 186–202, 465–473, 605–614, 669–675
intertidal zone 595–604
intraspecific competition 406–415, 525–534
intraspecific variation 465–473, 732–739
Iris gracilipes 633–641
irradiance 113–123, 124–135, 252–262:
  competition 155–164, 642–646
  and growth 145–154, 474–480
  seasonal 722–731

juvenile-phase invasive responses 103–112

knock-down resistance 289–296

Lacerta agilis 443–450
lactation 325–334
lakes 754–762
Lampornis amethystinus 78–84
landhoppers 435–442
leaf 3–12, 378–387, 490–496, 660–668:
  area ratios 113–123, 378–387, 804–811
  azimuth 124–135
  lifespan 351–359, 740–747, 748–753
  mass 113–123, 497–505, 804–811
  physiology 252–262, 378–387, 423–434
  pigment 474–480
  specific leaf area 113–123, 351–359, 688–695, 804–811
  traits 688–695
Licuala arbuscula 660–668
life history 29–35, 274–281:
  traits 165–174, 310–317, 569–574
light 660–668, 798–803:
  see also irradiance
Liriodendron tulipifera 263–273
liver and ventricle mass 60–69
lizards 186–202, 443–450
locomotion 186–202
locomotor speed 561–568
Lolium perenne 605–614
longevity 396–405, 712–721, 748–753
Lucilia sericata 712–721
lutein 297–303

marine benthic ecology 595–604
mass 60–69, 274–281, 325–334
maternal effects 13–23, 624–632, 772–781
maternal mass 325–334
maturation 310–317
Mediterranean 124–135, 186–202, 669–675
MeJA-induced response 406–415
Melanocallis caryaefoliae 525–534
melezitose 544–550
mice 575–583, 791–797
microarthropods 515–524, 647–653
microclimate 243–251
microcosms 51–59
microsite light environment 798–803
migration 310–317, 465–473, 584–594
millipedes 51–59
minnows 304–309
Mirafra erythroclamys 175–185
mistletoe 782–790
mites 78–84, 515–524
models 155–164, 233–242, 615–623, 701–711

© 2001 British Ecological Society,

Monellia caryella 525–534
Monoctonus paulensis 335–343
monospecific stand 642–646
morphometrics 203–210
mortality 24–28, 712–721, 748–753
moth 217–221
moult 29–35
Moussonia deppeana 78–84
Mus musculus 791–797
mutualism 544–550
mycorrhizal fungi 451–457

Narthecium asiaticum 396–405
nectar 78–84, 782–790
neighbour effect 642–646
nematodes 506–514, 569–574
Nephila clavipes 24–28
nest-site selection 772–781
net ecosystem carbon exchange 388–395
New Zealand 233–242, 497–505
Nicotiana attenuata 406–415
nitrogen 3–12, 155–164, 378–387, 605–614:
  allocation 406–415
  deficiency 165–174, 451–457
  storage 370–377, 535–543
  use 423–434
non-linear model 615–623
Nothofagus fusca 490–496
nursery streams 310–317
nutrient constraints 378–387, 423–434
nutritional deficiency 306–309, 318–324, 351–359, 791–797

offspring 335–343, 406–415, 561–568:
  survival 211–216, 443–450, 772–781
old-field succession 506–514
optimal temperature 701–711
Oreina elongata 318–324
osmoregulation 647–653
Osyris 669–675
overwintering reserves 370–377
oviposition 318–324

palms 252–262, 660–668, 804–811
Panicum dichotomiflorum 103–112
parasitoid wasps 217–221, 335–343
parent–offspring conflict 13–23
parental age 211–216
Passer montanus 29–35
patchiness 155–164
pelvic aperture 70–77
perch 754–762
perennial species 370–377, 406–415, 605–614:
  Australian 423–434
  grasses 155–164, 605–614, 748–753
  herbs 243–251, 633–641, 669–675, 740–747
Peromyscus maniculatus 575–583
petiole 660–668
Phalacrocorax aristotelis 211–216
phenology 722–731
phenotype plasticity 36–42, 96–102, 124–135, 740–747
phenotype variation 624–632
phenotypic flexibility 465–473
Phillyrea 669–675
Phoca vitulina 325–334
phosphorus 124–135, 165–174, 378–387
photochemical efficiency 124–135
photon flux density 798–803
photosynthesis 3–12, 243–251, 252–262, 263–273, 497–505, 740–747
photosynthetic adaptation 722–731
photosynthetic enzymes 423–434
photosynthetic pigments 474–480
Phoxinus phoxinus 304–309
phylogenic information 186–202
pigments 297–303, 474–480, 732–739
pike 754–762
Pinus sylvestris 535–543
Pinus taeda 344–350
Plagiothecium denticulatum 451–457
plankton, herbivorous 165–174
plant competition 642–646
plant distribution 222–232, 243–251
plant functional types 676–687
plant strategies 423–434
Plantago lanceolata 605–614
plasma metabolites 584–594
plasticity 24–28, 406–415, 481–489
plumage variation 732–739
plume loading 274–281
Poa pratensis 605–614
pollen tubes 782–790
pollination 78–84, 396–405, 551–558, 782–790:
  geitonogamous 633–641
polyols 647–653
population density 96–102, 575–583
population dynamics 712–721
Populus spp. 544–550
posthatching growth 624–632, 772–781
predation risk 29–35, 304–309, 318–324
preference–performance relationship 318–324
prey availability 36–42, 754–762
protandry 78–84
Protaphorura armata 647–653
protein 13–23, 370–377, 497–505

Quercus coccifera 124–135
quitting-harvest rate 575–583

Ratkowsky growth model 701–711
red to far-red light ratio 798–803
refuge 304–309
rehydration procedure 688–695
resistance 569–574
relative growth rate 113–123, 145–154, 274–281, 804–811
reproductive investment 13–23, 783–790
reproductive strategy 70–77
reptiles 36–42, 282–288, 561–568
residency 310–317
resistance 217–221, 406–415, 569–574
resource 3–12, 85–95, 615–623:
  depletion 165–174, 740–747
  dynamics 676–687
respiration 378–387, 423–434, 497–505, 791–797
respirometry 417–422, 458–464
resting metabolic rate 458–464
restoration 506–514
rhizodeposition 43–50
rice paddy 798–803
risk 24–28, 304–309, 318–324, 654–659
roach 754–762
root 3–12, 360–369, 370–377, 605–614:
  and leaf turnover 748–753
  zone water deficit 233–242
Rubia 669–675
Rubisco 3–12, 722–731
Rubus 669–675
Rumex acetosa 370–377
ruminants 569–574

Salmo salar 13–23, 701–711
Salmo trutta 310–317, 654–659
salt marsh 360–369
salt spray 435–442
saplings 113–123, 263–273, 535–543
savanna trees 222–232
scarification of seed coat 669–675
Scatophaga stercoraria 96–102
scramble competition 615–623

© 2001 British Ecological Society,

seals 325–334
seasonal change 388–395, 722–731
seasonal variation 465–473
seed 78–84, 145–154, 344–350, 669–674, 763–771
seedlings 145–154, 274–281, 344–350, 490–496, 763–771
self-shading 660–668
senescence 3–12
Setaria geniculata 103–112
settlement 595–604
sexual differences 561–568
sexual dimorphism 203–210, 211–216
sexual selection 297–303, 732–739
shade tolerance 113–123, 263–273
shrubs 145–154, 274–281, 451–457, 669–675, 782–790
size 96–102, 203–210, 561–568, 569–574:
  clutch 335–343
  eggs 13–23
  seed 763–771
  turtles 70–77, 481–489, 624–632, 772–781
smolting 310–317
snakes 36–42, 561–568
soil 43–50, 51–59, 490–496, 506–514, 748–753:
  and atmosphere 233–242
Sorbus aucuparia 535–543
spatial arrangement 633–641
spatial model 155–164
species composition 51–59, 605–614
species distribution 222–232, 233–242
species richness 51–59, 145–154
specific leaf area 113–123, 351–359, 688–695, 740–747, 804–811
spiders 24–28
Spinotarsus sp. 51–59
spring ephemeral 722–731
sprint performance 186–202
standard metabolic rate 624–632
statistical models 233–242
Sternotherus odoratus 70–77
stomatal limitation 222–232
stomatal regulation 490–496
stopover sites 584–594
stress tolerance 222–232
sugars 544–550, 647–653
sun/shade acclimation 474–480
sunflecks 263–273
Sylvia melanocephala 669–675

Taeniopygia guttata 458–464
taproot 370–377
Taraxacum officinale 605–614
Teladorsagia circumcincta circumcincta 569–574
temperature 217–221, 282–288, 351–359, 378–387
teratocytes 335–343
testis size 96–102
Thamnophis marcianus 36–42
thermal adaptation 701–711
thermal variance 561–568
thermoregulation 282–288
thermotolerance 217–221, 289–296
Tiarella cordifolia 722–731
tissue density 351–359
topological index 360–369
trade-offs 96–102, 145–154, 443–450
trait correlations 85–95
trees 145–154, 344–350, 551–558, 798–803:
  Australian 222–232, 351–359, 423–434
  boreal 451–457, 575–583
  cloud forest 78–84
  Mediterranean 124–135
  New Zealand 232–242, 490–496
  temperate 263–273, 425–434, 497–505, 535–543, 544–550, 804–811
  tropical rain 113–123, 378–387, 490–496, 763–771
  tropical transitional 388–395
  understorey 263–273, 798–803:
    palm 252–262, 660–668, 804–811
Trichogramma carverae 217–221
Trifolium repens 103–112, 605–614
Tristerix corymbosus 782–790
trophic complexity 506–514
Tropicoseius sp. nov. 78–84
Tropidonophis mairii 561–568
Turdus merula 669–675
turnover 155–164
turtles 70–77, 481–489, 624–632, 772–781

understorey 252–262, 263–273, 660–668, 798–803, 804–811

Vaccinium myrtillus 451–457
Valdensia heterodoxa 451–457
variance components 584–594
variation, reserves 481–489
variation, metabolic rate 465–473
vegetational composition 451–457
Viola pubescens 722–731
Virola michelii 490–496
volatile compounds 551–558

wasps 217–221, 335–343
water 175–185, 490–496, 740–747:
  use 222–232, 423–434, 605–614, 647–653
weather 282–288
wheat 615–623
wing wear 417–422
woodland 145–154, 351–359

Xanthium canadense 642–646
xanthophyll cycle 474–480

zygote dispersal 595–604

© 2001 British Ecological Society,